

DAFTAR PUSTAKA

- Abdollahi A, Shoar S, Nayyeri F, and Shariat M. 2012. Diagnostic value of simultaneous measurement of procalcitonin, IL-6, and hs-CRP in prediction of early-onset neonatal sepsis. *Mediterranean Journal of Hematology and Infectious Diseases*. 4 (1):e2012028. Doi:10.4084/mjhid.028.
- Alphonsus C S and Rodseth R N. 2014. The endothelial glycocalyx: a review of the vascular barrier. *Anaesthesia*. 69(7):777-784 doi:10.1111/anna.12661.
- Amersfoort E S V, Berkel T J C V, and Kuiper J. 2003. Receptors, Mediators, and Mechanisms Involved in Bacterial Sepsis and Septic Shock. *Clinical Microbiology Reviews*. Jul; 16(3): 379-414.
- Anas A A, Wiersinga W J, deVos A F, and Van Der Poll T. 2010. Recent Insight in to the Pathogenesis of Bacterial Sepsis. *The Netherlands Journal of Medicine*. Apr; 68(4): 147-52.
- Annecke T, Fischer J, Hartmann H, Tschoep J, Rehm M, Conzen P, Sommerhoff C P, and Becker B F. 2011. Shedding of the coronary endothelial glycocalyx: effects of hypoxia/reoxygenation vs ischaemia/reperfusion. *British Journal of Anaesthesia*. 107(5):679-686 doi:10.1093/bja/aer269.
- Arulselvan P, Tan W S, Gothai S, Muniandy K, Fakurazi S, Esa N M, Abdullah A. Alarfaj, and Kumar S S. 2016. Anti-Inflammatory Potential of Ethyl Acetate Fraction of *Moringa oleifera* in Downregulating the NF- κ B Signaling Pathway in Lipopolysaccharide-Stimulated Macrophages in Molecules. www.mdpi.com/journal/molecules. 21, 1452; doi:10.3390/molecules21111452.
- Atawodi S E, Atawodi J C, Idakwo G A, Pfundstein B, Haubner R, Wurtele G, Bartsch H, and Owen R W. 2010. Evaluation of the polyphenol content and antioxidant properties of methanol extracts of the leaves, stem, and root barks of *Moringa oleifera* Lam. *Journal of Medicinal Food*. Doi: 10.1098/jmf.2009.0057.
- Aygün F O, Akçam F Z, Kaya O, Ceyhan B M, and Sütçü R. 2012. Caffeic acid phenethyl ester modulates gentamicin-induced oxidative nephrotoxicity in kidney of rats. *Biological Trace Element Research*. 145(2):211-6.
- Baratawidjaja K G and Iris R. 2009. *Imunologi Dasar*. Edisi VIII. FK Universitas Indonesia.
- Becker B F, Chappell D, Bruegger D, Annecke T, and Jacob M. 2010. Therapeutic strategies targeting the endothelial glycocalyx: acute deficits, but great potential. *Cardiovascular Research*. 87(2):300-310 doi: 10.1093/cvr/cvq137
- Ben-Izhak O, Kaplan-Cohen V, Ilan N, Gan S, Vlodavsky I, and Nagler R. 2006. Heparanase expression in malignant salivary gland tumors inversely with long-term survival. *Neoplasia*. 8(10):879-884 doi: 10.1593/neo.06382.
- Bone R C, Balk R A, Cerra F B, Dellinger R P, Fein A M, Knaus W A, Schein R M, and Sibbald W J. 1992. Definitions for sepsis and organ failure and guidelines for

the use of innovative therapies in sepsis. The ACCP/SCCM Consensus Conference Committee. *American College of Chest Physicians/Society of Critical Care Medicine Chest*. Jun;101(6):1644-55. DOI:10.1378/chest.101.6.1644.

Bouza C, Cuadrado L T, Parkinson S Z, and Blanco A J M. 2014. Epidemiology and recent trends of severe sepsis in Spain: a nationwide population-based analysis (2006-2011). *BMC Infectious Diseases*. 14: 717. Research article.

Broekhuizen L N, Mooij H L, Kastelein J J P, Stroes E S G, Vink H, and Nieuwdorp M. 2009. Endothelial glycocalyx as potential diagnostic and therapeutic target in cardiovascular disease. *Current Opinion in Lipidology*. 20(1):77-62 doi: 10.1097/MOL.0b013e328321b587.

Burke-Gaff A and Evans T W. 2012. Lest we forget the endothelial glycocalyx in sepsis. *Burke-Gaffney and Evans Critical Care*. 16:121.

Chelazzi C, Villa G, Marcinelli P, De Gaudio A R, and Adembri C. 2015. Glicocalyx and sepsis-induced alterations in vascular permeability. *Critical Care*. 19(1, article 26) doi: 10.1186/s13054-015-0741-z.

Chen G, Wang D, Vikramadithyan R, Yagyu H, Saxena U, Pillarisetti S, and Goldberg I J. 2004. Inflammatory cytokines and fatty acids regulate endothelial cell heparanase expression. *Biochemistry*. 43(17):4971-4977 doi: 10.1021/bi0356552.

Cho S Y, Park S J, Kwon M J, Jeong T S, Bok S H, Choi S H, Choi W Y, Jeong W I, Ryu S Y, Do S H, Lee C S, Song J C, and Jeong K S. 2003. Quercetin suppresses pro-inflammatory cytokines production through MAP kinases and NF-KappaB pathway in lipopolysaccharide-stimulated macrophage. *Molecular and Cellular Biochemistry*. 243:153-160.

Choi Y H, Kim G Y, and Lee H H. 2014. Anti-inflammatory effects of cordycepin in lipopolysaccharide-stimulated RAW264.7 macrophages through toll-like receptor 4-mediated suppression of mitogen-activated protein kinase and NF- κ B signaling pathways. *Drug, Design, Development and Therapy*. 8, 1941-1953.

Choi Y Y, Kim M H, Han J M, Hong J, Lee T H, Kim S H, and Yang W M. 2014. The anti-inflammatory potential of cortex phellodendron in vivo and in vitro: Down-regulation of no and inos through suppression of NF- κ B and MAPK activation. *International Immunopharmacology*. 19, 214-220.

Cruz K and Dellinger P R. 2003. Diagnosis and Source of Sepsis: The Utility of Clinical Findings. *The Sepsis Text pp. 11-28.*

Cui N, Wang H, Long Y, Su L, and Liu D. 2015. Dexamethasone Suppressed LPS-Induced Matrix Metalloproteinase and Its Effect on Endothelial Glycocalyx Shedding. *Mediators of Inflammation*. 912726.

Cummings C J, Sessler C N, Beall L D, Fisher B J, Best A M, and Fowler A A. 1997. Soluble e-Selectin levels in sepsis and critical illness. Corelations with infections

and hemodynamic dysfunction. *American Journal of Respiratory and Critical Care Medicine.* aup; 156 (2 Pt 1): 431-7.

Dantes N. 2012. *Metode Penelitian.* Yogyakarta: Penerbit Andi.

Doi K, Leelahanichkul A, Peter S T, Yuen, and Robert A S. 2009. Animal models of sepsis and sepsis-induced kidney injury. *The Journal of Clinical Investigation.* 119(10):2868-2878. <https://doi.org/10.1172/JCI39421>.

Doreen E W, Swan R Z, Chung C, and Ayala A. 2007. The Apoptotic Pathway as a Therapeutic Target in Sepsis. *Current Drug Targets.* 8(4): 493-500.

Dutordoir M R and Bates D A V. 2016. Activation of Apoptosis Signalling Pathways by Reactive Oxygen Species. *Biochmica et Biophysica Acta.* [Doi.org/10.1016/j.bbamcr.2016.09.012](https://doi.org/10.1016/j.bbamcr.2016.09.012).

Elmore S. 2007. Apoptosis: A Review of Programmed Cell Death. *Toxicologic Pathology.* 35(4): 495–516.

Endang T and Sukma D. 2016. Fraksi Metanol Daun Kelor Menurunkan Kadar TNF- α dan IL-6 Serum, serta MDA Kolon Tikus yang Diinduksi DMBA. *Jurnal Kedokteran Brawijaya.* Vol. 29, No. 1.

Eroschenko V P. 2008. Circulatory system. In: Eroschenko, VP, editor. DiFiore's Atlas of Histology with functional correlation. 11th edition. Philadelphia: Lippincott Williams & Wilkins. p.313-29.

Exline M C and Crouser E D. 2014. Mitochondrial Mechanisms of Sepsis-Induced Organ Failure. *NIH Public Acces.* 13:5030-5041.

Federer W T. 1995. Experimental design: theory and application. *Journal Osborne.* Macmillan, New York.

Gill S E, Rohan M, and Mehta S. 2015. Role of pulmonary microvascular endothelial cell apoptosis in murine sepsis-induced lung injury in vivo. *Respiratory Research.* 16(1, article 109) doi: 10.1186/s12931-015-0266-7.

Goldberg R, Meirovitz A, Hirshoren N, Bulvik R, Binder A, Rubinstein A M, and Elkin M. 2013. Versatile role of heparanase in inflammation. *Matrix Biology.* 32(5):234-240 doi: 10.1016/j.matbio.

Goodall K J, Poon I K H, Phipps S, and Hulett M D. 2014. Soluble heparan sulfate fragments generated by heparanase trigger the release of pro-inflammatory cytokines through TLR-4. *PLoS ONE.* 9(10) doi: 10.1371/journal.pone.0109596.e109596.


Guha M and Mackman N. 2001. LPS induction of gene expression in human monocytes. *Cell Signal.* 13, 85-94.

Guntur H A, Diding H P, Pohan H T, and Widodo D. 2011. Effect of low-dose steroid on NF- κ B and caspase-3 intestinal expression in a sepsis mouse model. *Critical care*. Vol 15, Suppl 3.

Guntur H A. 2009. Development of Immunopathobiogenesis on SIRS-Sepsis. *The Indonesian Biomedical Journal*. Vol.1, No.1, April 2009 p.32-9.

Heo S J, Jang J, Ye B R, Kim M S, Yoon W J, Oh C, Kang D H, Lee J H, Kang M C, Jeon Y J, Kang S M, Kim D, and Kim K N. 2014. Chromone suppresses the activation of inflammatory mediators in lipopolysaccharide-stimulated RAW264.7 cells. *Food and Chemical Toxicology*. 67, 169-175.

Ibrahim K E, Al-Mutary M G, Bakhet A O, and Khan H A. 2018. Histopathology of the Liver, Kidney; Spleen of Mice Exposed to Gold Nanoparticles. *Molecules*. 23, 1848; doi:10.3390/molecules23081848.



Ince C, Mayeux P R, Nguyen T, Gomez H, Kellum J A, Ospina-Tascón G A, Hernandez G, Murray P, and De Backer D. 2016. The Endothelium In Sepsis Shock. *ADQI XIV Workgroup*. 45(3): 259–270.

Kapplan M J and Radic M. 2012. Neutrophil extracellular traps: double-edged swords of innate immunity: *Journal of Immunology*. 189(6):2689-2695 doi: 10.4049/jimmunol.1201719.

Kenneth T. 2012. *Online Texbook of Bacteriology*. www.texbookofbacteriology.net.

Konrad R, Michael B, Niels C, Riedemann, and Christiane S Hartog. 2012. New Approaches to Sepsis: Molecular Diagnostic and Biomarkers. *Volume 25 Number 4 Clinical Microbiology Reviews* p. 609-34.

Kou X, Li B, Olayanju J B, Drake J M, and Chen N. 2018. Nutraceutical or Pharmacological Potential of *Moringa Oleifera Lam*. www.mdpi.com/journal/nutrients. 343; doi:10.3390/nu10030343.

Kovach M A and Standiford T J. 2012. The function of neutrophils in sepsis. *Current Opinion in Infectious Diseases*. 25(3):321-327 doi: 10.1097/QCO.0b013e3283528c9b.

Kranidioti H, Orfanos S E, and Vakil I. 2009. Angiopoietin-2 is increased in septic shock: evidence for the existence of a circulating factor stimulating its release from human monocytes. *Immunology Letters*. 125(1):65-71 doi 10.1016/j.imlet.2009.06.006.

Kusnul Z and Munir Z. 2012. Efek Pemberian Jus Menthimun Terhadap Penurunan Tekanan Darah. <http://repository.usu.ac.id/bitstream/123456789/26951/4/Chapter%20I2.pdf>

Lawrence T. 2009. IKK α in the regulation of inflammation and adaptive immunity. *Biochemical Society Transactions*. 35:270–272.

Ley K. 1996. Molecular mechanisms of leukocyte recruitment in the inflammatory process. *Cardiovaskular Research*. 32(4):733-742 doi: 10.1016/0008-6363(96)00066-1.

Ling J and Kumar R. 2012. Crosstalk between NF κ B and glucocorticoid signaling: A potential target of breast cancer therapy. *Cancer Letters*. 322, 119-126.

Liu C M, Sun Y Z, Sun J M, Ma J Q, and Cheng C. 2012. Protective role of quercetin against lead-induced inflammatory response in rat kidney through the ROS-mediated MAPKs and NF- κ B pathway. *Biochmica et Biophysica Acta*. 1820(10): 1693-703.

Lygizos M I, Yang Y, and Altman C J. 2013. Heparanase mediates renal dysfunction during early sepsis in mice. *Physiological Reports*. 1(6) doi: 10.1002/phy2. 153. e 00153.

Mak T W and Saunders M E. 2004. The Immune Response Basic and Clinical Principles. *Shock*. Hal: 206-208.

Mantzarlis K, Tsolaki V, and Zakynthinos E. 2017. Role of Oxidative Stress and Mitochondrial Dysfunction in Sepsis and Potential Therapies. *Hindawi Oxidative Medicine and Cellular Longevity*. <https://doi.org/10.1155/2017/5985209>.

Martin L, De Santis R, and Koczera P. 2015. The synthetic antimicrobial peptide 19-2.5 interacts with heparanase and heparan sulfate in murine and human sepsis. *PLoS ONE*. 10(11) doi: 10.1371/journal.pone. 0143583. e 0143583.

Martin L, Koczera P, Zechendorf E, and Schuerholz T. 2016. The Endothelial Glycocalyx: New Diagnostic and Therapeutic Approaches in Sepsis. *Biomed Research International*. 3758278.

Martin L, Peters C, and Schmitz S. 2015. Soluble heparan sulfate in serum of septic shock patients induces mitochondrial dysfunction in murine cardiomyocytes. *Shock*. 44(6):569-577. doi: 10:1097/SHK.0000000000000462.

Martin L, Schmitz S, and De Santis R. 2015. Peptide 19-2.5 inhibits heparan sulfate-triggered inflammation in murine cardiomyocytes stimulated with human sepsis serum. *PLoS ONE*. 10(5) doi: 10.1371/journal.pone. 0127584. e 0127584.

Mercurio F and Manning A M. 1999. Multiple signals converging on NF- κ B. *Current Opinion in Cell Biology*. 11, 226-232.

Michael D, Howell, and Andrew M. 2017. Davis Management of Sepsis and Septic Shock. *JAMA Clinical Guidelines Synopsis*. 317(8): 847-848.

Namas R, Zamora R, Gary M D, Doyle J, Thomas E, Frank J, Jacono M D, Androulakis I P, Nieman G F, Chang S, Billiar T R, Kellum, Angus D C, and Vodovotz Y. 2012. Sepsis: something old, something new, and a system view. *The Journal of Critical Care*. 27:314.e1-314.e11.

Nelson A, Berkstedt I, and Bodelsson M. 2014. Circulating glycosaminoglycan species in septic shock. *Acta Anaesthesiologica Scandinavica*. 58(1):36-43 doi: 10.1111/aas.12223.

Nieuwdorp M, Meuwese M C, Vink H, Hoekstra J B L, Kastelein J J P, and Stroes E S G. 2005. The endothelial glycocalyx: a potential barrier between health and vascular disease. *Current Opinion in Lipidology*. 16(5):507-511 doi: 10.1097/01.mol.0000181325.08926.9c.

Niu X, Xing W, Li W, Fan T, Hu H, and Li Y. 2012. Isofraxidin exhibited anti-inflammatory effects in vivo and inhibited TNF- α production in LPS-induced mouse peritoneal macrophages in vitro via the MAPK pathway. *International Immunopharmacol* 14, 164-171.

Northeastern University. 2012. Qualitative research methods overview. <http://www.ccs.neu.edu/course/is4800sp12/resources/qualmethods.pdf>.

Osterbur K, Mann F A, Kuroki K, and Decleu A. 2014. Multiple Organ Dysfunction Syndrome in Humans and Animals. *Journal of Veterinary Internal Medicine*. Jul-Aug; 28(4): 1141-1151.

Ouédraogo M, Lamien-Sanou A, Ramde N, Ouédraogo A S, Moussa O, Zongo S P, Goumbri O, Duez P, and Guissou P I. 2013. Protective effect of *Moringa oleifera* leaves against gentamicin-induced nephrotoxicity in rabbits. *Science Direct*. <https://doi.org/10.1016/j.etp.2011.11.006>.

Perkins N D. 2000. The REL/ NF- κ B family: Friend and foe. *Trends in Biochemical Science*. 25, 434-440.

Poll V D T, De Boer J D D, and Levi M. 2011. The effect of inflammation on coagulation and vice versa. *Current opinion in Infectious Diseases*. 24(3):273-278 doi: 10.1097/QCO.0b013e328344c078.

Putri A, Setyawati T, and Lintin G. 2016. Efek Antibakteri Ekstrak Daun Kelor (*Moringa Oleifera*) Terhadap Pertumbuhan Bakteri *Salmonella Thypi*. xvi, 42 hlm.

Putri O H, Armalina D, Mundhofir F E P, Ismail A, and Pawitra I. 2018. Pengaruh Pemberian Fraksi Daun Kelor (*Moringa Oleifera*) Dosis Bertingkat pada Gambaran Mikroskopis Hepar Tikus Wistar yang Diinduksi Formalin. *Jurnal Kedokteran Diponegoro*. 7 (2): 2540-8844.

Rajapakse N, Kim M M, Mendis E, and Kim S K. 2008. Inhibition of inducible nitric oxide synthase and cyclooxygenase-2 in lipopolysaccharide-stimulated RAW264.7 cells by carboxybutyrylated glucosamine takes place via down-regulation of mitogen-activated protein kinase-mediated nuclear factor- κ B signaling. *Immunology*. 123, 348-357.

Rehm M, Bruegger D, Christ F, Christ F, Conzen P, Thiel M, Jacob M, Chappell D, Stoeckelhuber M, Welsch U, Reichart B, Peter K, and Becker B F. 2007. Shedding of the endothelial glycocalyx in patients undergoing major vascular surgery with global and regional ischemia. *Circulation*. 116(17):1896-1906 doi: 10.1161/CIRCULATIONAHA.106.684852.

Remick D G. 2007. Pathophysiology of Sepsis. *The American Journal of Pathology*. 170 (5): 1435-44. Doi: 10.2353/ajpath.2007.060872.

Sallisalmi M, Tenhunen J, Yang R, Oksala N, and Pettilä V. 2012. Vascular adhesion protein-1 and syndecan-1 in septic shock. *Acta Anaesthesiologica Scandinavica*. 56(3):316-322 doi: 10.1111/j.1399-6576.2011.02578.x.

Sarrazin S, Lyon M, Deakin J A, Guerrini M, Lassalle P, Delehedde M, and Lortat-Jacob H. 2010. Characterization and binding activity of the chondroitin/dermatan sulfate chain from Endocan, a soluble endothelial proteoglycan. *Glycobiology*. 20(11):1380-1388 doi 10.1093/glycob/cwq100.

Scherpereel A, Depontieu F, Grigoriu B, Cavestri B, Tsicopoulos A, Gentina T, Jourdain M, Pugin J, Tonnel A B, and Lassalle P. 2016. Endocan, a new endothelial marker in human sepsis. *Critical Care Medicine*. 34(2):532-537 doi: 10.1097/01.CCM.0000198525.82124.74.

Schmidt E P, Yang Y, Janssen W J, Gandjeva A, Perez M J, Barthel L, Zemans R L, Bowman J C, Koyanagi D E, Yunt Z X, Smith L P, Cheng S S, Overdier K H, Thompson K R, Geraci M W, Douglas I S, Pearse D B, and Tuder R M. 2012. The pulmonary endothelial glycocalyx regulates neutrophil adhesion and lung injury during experimental sepsis. *Nature Medicine*. 18(8):1217-1223 doi: 10.1038/nm.2843.

Shafat I, Ilan N, Zoabi S, Vlodavsky I, and Nakhoul F. 2011. Heparanase levels are elevated in the urine and plasma of type 2 diabetes patients and associate with blood glucose levels. *PLoS ONE*. 6(2) doi:10.1371/journal.pone.0017312. e17312.

Shaker O, Zahra A, Sayed A, Refaat A, Hegazy G, El-Khaiat Z, El-Hindawi K, and Ay-El Deen M. 2013. Role of ICAM-1 and E-selectin gene polymorphisms in pathogenesis of PAOD in Egyptian patients. *Vascular Health and Risk Management*. 2013:6.

Siner J M, Bhandari V, Engle K M, Elias J A, and Siegel M D. 2009. Elevated serum angiopoietin 2 levels are associated with increased mortality in sepsis. *Shock*. 31(4):348-353 doi: 10.1097/shk.0b013e318188bd06.

Soldato W, Swan R Z, Chung S C, and Ayala A. 2007. The Apoptotic pathway as therapeutic target in sepsis. *Current Drug Targets*. 8: 493-500.

Song J W, Zullo J, Liphhardt M, Dragovich M, Zhang F X, Fu B, and Goligoesky M S. 2017. Endothelial glycocalyx – the battleground for complications of sepsis and kidney injury. *Nephrology Dialysis Transplantation*. (4) 22: 1- 9.

Steppan J, Hofer S, Funke B, Brenner T, Henrich M, Martin E, Weitz J, Hofmann U, and Weigand M A. 2011. Sepsis and major abdominal surgery lead to flaking of the endothelial glycocalix. *The Journal of Surgical Research*. 165(1):136-141 doi: 10.1016/j.jss.2009.04.034.

Sukamto I G L, Purwanto B, and Suroto M A. 2018. The Optimal Dose of Intraperitoneal Head Shock Protein (HSP70) to Prevent Death in Sepsis Mice Model With Multiple

Organ Dysfunction Syndrome. *Mid Internal Conference on Public Health.* Surakarta. Doi: <https://doi.org/10.26911/Mid.ICPH.2018.05.21>

Sulistyawati R, Nurani L H, Hidayati S, Mursyidi A, and Mustofa. 2017. Standarisasi Kualitas Fraksi Etil Asetat Daun Kelor (*Moringa oleifera* Lamk.). *University Research Colloquium.* Universitas Muhammadiyah Magelang.

Sumandjar T, Purwanto B, Riswanto, and Rahman A. 2019. Effects of Ethyl Acetate Fraction in *Moringa oleifera* Leaves on liver and kidney dysfunction in sepsis mice (Study of NF-κB expression in the aorta and renal proximal tubules, SGOT, SGPT, urea and creatinine). *Balimedicaljournal, In press.*

Surviving Sepsis Campaign: International Guidelines for Management of Sepsis and Septic Shock. *JAMA Clinical Guidelines Synopsis.* 2016;315(8):801-810. doi:10.1001/jama.2016.0287.

Tan, W S, Aruselvan P, Karthivashan G, and Fakurazi S. 2015. *Moringa oleifera* flower extract suppresses the activation of inflammatory mediators in lipopolysaccharide-stimulated RAW264.7 macrophages via NF-κB pathway. *Mediators of Inflammation.* 720171.

Taufiqqurrohman M A. 2008. *Pengantar Metodologi Penelitian Untuk Ilmu Kesehatan.* Surakarta: LPP UNS dan UNS Press.

Ushiyama A, Kataoka H, and Iijima T. 2016. Glycocalyx and its involvement in clinical pathophysiologies. *Journal of Intensive Care.* (4):59.

Vlodavsky I, Beckhove P, and Lerner I. 2012. Significance of heparanase in cancer and inflammation. *Cancer Microenvironment.* 5(2):115-132 doi: 10.1007/s12307-011-0082-7.

Wang X and Lau H. 2006. Prostaglandin E₂ potentiates the immunologically stimulated histamine release from human peripheral blood-derived mast cells through EP1 / EP3 receptors. *Allergy.* 61, 503-506.

Waterman C, Cheng, Diana M, Rojas-Silva P, Poulev A, Dreifus J, Lila M A, and Raskin I. 2014. Stable, water extractable isothiocyanates from *Moringa Oleifera* leaves attenuate inflammation in vitro. 103:114-122. doi: 10.1016/j.phytochem.2014.03.028.

Waterman C, Rojas-Silva P, Tumer T B, Kuhn P, Richard A J, Wicks S, Stephens J M, Wang Z, Mynatt R, and Cefalu W. 2015. Isothiocyanate-rich *Moringa oleifera* extract reduces weight gain, insulin resistance, and hepatic gluconeogenesis in mice. *Molecular Nutrition & Food Research.* 59, 1013-1024.

Weber S U, Schewe J C, Lehmann L E, Muller S, Book M, Klaschick S, Hoeft A, and Stuber F. 2008. Induction of Bim and Bid gene expression during accelerated apoptosis in severe sepsis. *Critical Care.* 12: R 128 doi:10.1186/cc7088.

Wei W C, Lin S Y, Chen Y J, Wen C C, Huang C Y, Palanisamy A, Yang N S, and Sheu J H. 2011. Topical application of marine briarane -type diterpenes effectively inhibits 12-O- tetradecanoylphorbol-13-acetate-induced inflammation and dermatitis in murine skin. *Journal of Biomedical Science*. 18, 94.

Weinbaum S, Tarbell J M, and Damiano E R. 2007. The structure and function of the endothelial glycocalyx layer. *Annual Review of Biomedical Engineering*. 9:121-167 doi: 10.1146/annurev.bioeng.9.060906.151959.

Williams S H, Che X, Paulick A, Guo C, Lee B, Muller D, Uhlemann A C, Lowy F D, Corrigan R M, and Lipkin W I. 2018. New York City House Mice (*Mus musculus*) as Potential Reservoirs for Pathogenic Bacteria and Antimicrobial Resistance Determinants. *American Society For Microbiology*. Mar-Apr; 9(2): e00624-18.

Wulan A H, Sholeh S N, and Wigati D. 2017. Pengaruh Pemberian Fraksi Etil Asetat Daun Kelor (*Moringa Oleifera Lam.*) Tehadap Gambaran Histopathology dan Kadar SGPT dan SGOT pada Tikus Jantan Galur Wistar yang Induksi Monosodium Glutamat. *Media Farmasi Indonesia*; 14 (1).

Xing K, Murthy S, Liles W C, and Singh J M. 2012. Clinical utility of biomarkers of endothelial activation in sepsis-a systematic review. *Critical Care*. doi: 10.1186/cc11145.

Yang Y, Haege S M, Suflita M A, Zhang F, Dailey K L, James F, Colbert, Ford J A, Picon M A, Stearman R S, Lin L, Liu X, Han X, Linhardt R J, and Schmidt E P. 2017. Fibroblast Growth Factor Signaling Mediates Pulmonary Endothelial Glycocalyx Reconstitution. *American Journal of Respiratory Cell and Molecular Biology*. Vol 56, Iss 6, pp 727–737.

Yeom M, Park J, Lim C, Sur B, Lee B, Han J J, Choi H D, Lee H, and Hahn D H. 2015. Glucosylceramide attenuates the inflammatory mediator expression in lipopolysaccharide-stimulated RAW264.7 cells. *Nutrition Research*. 35, 241-250.

Yuliani N N and Dienina D P. 2015. Uji Aktivitas Antioksidan Infusa Daun Kelor (*Moringa Oleifera*, Lamk) dengan Metode 1,1- Diphenyl-2-Picrylhydrazyl (DPPH). *Jurnal Info Kesehatan*. Vol. 14, Nomor 2.

Zhou J, Zhang HA, Lin Y, Liu HM, Cuy YM, Xu Y, Zhao N, Ma JM, Fan K, and Jiang CL. 2014. Protective effect of ginsenoside against acute renal failure via reduction of renal oxidative stress and enhanced expression of chat in the proximal convoluted tubule and ERK1/2 in the paraventricular nuclear. *Physiological Research*. 63(5):597-604.